

# **BPOC**

**Dr. C Baicus**

**[www.baicus.ro](http://www.baicus.ro)**

# Definitii

- BPOC:
  - Boala obstructiva pulmonara, progresiva si ireversibila
  - Modificari multiple pulmonare, incluzand parenchimul si caile aeriene mici
  - Efecte extra-pulmonare si comorbiditati importante, care influenteaza severitatea bolii
  - Declinul pe termen lung al functiei pulmonare

# Definitii


- Bronsita cronica:
  - Tuse & sputa pentru  $\geq 3$  luni/an, cel puțin 2 ani consecutivi
- Emfizemul:
  - Termen morfologic ce descrie distrugerea suprafetelor la nivelul carora au loc schimburile de gaze (alveole)

# Definitii

- Exacerbarea BPOC:
  - Aparitia acuta, in evolutia naturala a bolii, a unui episod caracterizat prin accentuarea dispneei bazale, a tusei si a expectoratiei

# Asthma and COPD


---



---

*Reproduced from the Global Initiative for Chronic Obstructive Pulmonary Disease, based on an April 1998 meeting of the National Heart, Lung, and Blood Institute and the World Health Organization.*

# Chronic obstructive pulmonary disease





# Incidentă/Prevalență

- A 4-a cauza de mortalitate în lume
- Prevalență 4-10% în lume
- Fumatori >15 ani: 47% la spirometrie

# Cauze / FR

- Fumatul (x1,5)
- Hiperresponsivitate cai aeriene
- Inhalarea de fum de tigara, prafuri/chimicale, poluanti
- Poluare, infectii repetate in copilărie, TBC pulmonara, astm (x10), lipsa dezvoltării intrauterine, subnutritie, statut social scazut
- Deficit de alpha-1 antitripsina (homo/heteroz)
- Biomasa (fum, combustibil) (x2,5)
- Marijuana (x3)










**FIGURE 2-8** Age-related rate of decline in lung function in various patient groups. *FEV<sub>1</sub>*, forced expiratory volume in 1 second. (Modified with permission from Snider GL, Saling LJ, Renard SI. Chronic bronchitis and emphysema. In Murray JF, Nadel JA, eds. Textbook of Respiratory Medicine. Philadelphia: WB Saunders, 1994:1342.)

# Fiziopatologie

- Obstructie cai aeriene / Hiperinflatie
  - ↓ **VEMS**, CV, **IT**(=VEMS/CV),
  - ↑ vol rezidual, capacitat pulm totala
- Respiratia
  - ↓ PaO<sub>2</sub> (VEMS<50%)
  - ↑ PaCO<sub>2</sub> (VEMS<25%)
  - ↑ TAP, CPC (VEMS<25%, PaO<sub>2</sub><55 mmHg)

# Morfopatologie





© Elsevier. Herring: Learning Radiology: Recognizing the Basics

Figure 14-11 Types of emphysema. Centriacinar (centrilobular) emphysema (A) features focal destruction limited to the respiratory bronchioles and the central portions of the acinus (closed white arrows). It is associated with cigarette smoking and is most severe in the upper lobes. Panacinar (panlobular) emphysema (B) involves the entire alveolus distal to the terminal bronchiole, is most severe in the lower lung zones and generally develops in patients with homozygous  $\alpha_1$ -antitrypsin deficiency. Paraseptal emphysema (C) is the least common form; it involves distal airway structures, alveolar ducts, and sacs; tends to be subpleural; and may cause pneumothorax.

# Manifestari clinice

- Simptomele principale
  - Tuse
  - Expectoratie
  - Dispnee

# Emfizem / Br cr

	Emfizem	Br cr
Definitie	Dilatatie/distrugere spatii aeriene (definitie morfologica)	Tuse productiva >3 luni/an, $\geq 2$ ani (definitie clinica)
Fiziopat	Afectat parenchimul Deficit paralel V/P Hipoxemie usoara	Afectate caile aeriene mici Defecte potrivire V/P Hipoxemie severa, hipercapnie, HTP, cord pulmonar
Manif clinice	Dispnee severa, constanta Tuse usoara	Dispnee intermitenta Tuse cu expectoratie
Ex fizic	<i>Pink puffer</i> Tahipnee, fara cianoza Murmur vezicular asurzit	<i>Blue bloater</i> Cianoza, obezitate, edeme Ronflante

# Diagnostic


- Clinic: istoric, obstructie bronsica (expir prelungit, sibilante), torace emfizematos, murmur vezicular asurzit
- Fumator
- Utilizare musculatura accesorie
- Spirometrie
- RX, CT

# Diagnostic

		Sn (%)	Sp (%)	LR+	LR-
h laring ≤ 4 cm*		36	90	3,6	0,7
s Harzer		4-27	<b>97-99</b>	<b>7,4</b>	NS
No matit cord		15	<b>99</b>	<b>11,8</b>	NS
Scor sunete resp**	≤ 9	23-46	<b>96-97</b>	<b>10,2</b>	
	≥ 16	<b>3-10</b>	33-34	<b>0,1</b>	
Subcrepit inspir		25-77	<b>97-98</b>	<b>14,6</b>	
Sibilant		13-56	86-99	2,8	0,8
T expir fortat	< 3 s	<b>8-10</b>	26-62	<b>0,2</b>	
	≥ 9 s	29-50	86-98	4,1	
2 din 3 (≥ 70 pach-an; spune ca are br cr/emfizem; ↓ murmur vez)		67	<b>97</b>	<b>25,7</b>	<b>0,3</b>

\* Sup cartilaj tiroidian – marg sup stern / \*\*0-4pt, 3 zone x2

**Dg  $\neq$**



© Elsevier. Herring: Learning Radiology: Recognizing the Basics


Figure 14-12 Emphysema. On conventional radiographs, the imaging findings of chronic obstructive pulmonary disease (COPD) are hyperinflation, including flattening of the diaphragm, especially on the lateral exposure (B) (open white arrows), increase in the retrosternal clear space (dotted white arrow), hyperlucency of the lungs with fewer than normal vascular markings (closed white arrows) (A), and prominence of the pulmonary arteries secondary to pulmonary arterial hypertension (open black arrows).

## Centrilobular emphysema



HRCT shows multiple small lucencies permeating the upper lobes. The wall of the emphysematous spaces is imperceptible.  
*Courtesy of Paul Stark, MD.*

## Panlobular emphysema



HRCT shows a paucity of vascular structures in both lower lobes, most evident in the anterior-basal segment of the right lower lobe.

*Courtesy of Paul Stark, MD.*



Harrison's

**FIGURE 254-4** Chest CT scan of a patient with COPD who underwent a left single-lung transplant. Note the reduced parenchymal

# Stadializare

Stage	Spirometry Findings
<b>Stadiul I - usor</b>	<b>VEMS/CV &lt; 0.7</b> <b>VEMS ≥ 80% prezis</b>
<b>Stadiul II – moderat</b>	<b>VEMS/CV &lt; 0.7</b> <b>VEMS ≥ 50% si &lt; 80%</b> <b>prezis</b>
<b>Stadiul III – sever</b>	<b>VEMS/CV &lt; 0.7</b> <b>VEMS ≥ 30% si &lt; 50%</b> <b>prezis</b>
<b>Stadiul IV - foarte sever</b>	<b>VEMS/CV &lt; 0.7</b> <b>VEMS &lt; 30% prezis sau</b> <b>VEMS &lt; 50% prezis plus IRC</b>

# Indice BODE

Variabila	Puncte indice BODE			
	0	1	2	3
VEMS (% prezis)	$\geq 65$	50-64	36-49	$\leq 35$
Dist in 6min (m)	$\geq 350$	250-349	150-249	$\leq 149$
Scala dispneei	0-1	2	3	4
IMC	$>21$	$\leq 21$		

**7-10: † 80% la 2 ani**


- **VEMS < 0,750 L: suprav 5 ani = 25%**
- **VEMS = 1L: suprav 5 ani = 50%**
- **↓ VEMS cu 50-75 mL/an (x2 N)**

# Tratamentul

- Obiective

# Tratamentul de fond

## Oprirea fumatului



# Tratamentul de fond

- Oprirea fumatului sta la baza trat
- Prima linie: bronhodilatator cu actiune lunga (beta-agonist sau anticolinergic).
- Stadiul III sau IV; exacerbari frecvente: combinatie LABA+cortizon (reduce exacerbarile).
- Exercitiul si reabilitarea imbunatatesc toleranta la exercitiu si reduc dispneea si oboseala.
- Oxigenoterapie ( 15-20 ore/zi) imbunatateste supravietuirea la pac cu obstructie severa ( $PaO_2 < 55$  mm Hg in repaus).

# Tratamentul de fond


- Oprirea fumatului sta la baza trat
- Vaccin antigripal (anual) si antipneumococic (la 5 ani).
- Bronhodilatatoare (beta2-adrenergice, anticolinergice)
- Cortizon inhalator (VEMS<50% + exacerb frecv)
- Aminofilina, teofilina: ↓ dispnee (putin)
- Mucolitice: ?

I: Mild

II: Moderate

III: Severe

IV: Very Severe



Harrison's